

9月15日 までは本年値、それ以降は平年値を使用

		収 穫 期																																																											
		8/15	8/16	8/17	8/18	8/19	8/20	8/21	8/22	8/23	8/24	8/25	8/26	8/27	8/28	8/29	8/30	8/31	9/1	9/2	9/3	9/4	9/5	9/6	9/7	9/8	9/9	9/10	9/11	9/12	9/13	9/14	9/15	9/16	9/17	9/18	9/19	9/20	9/21	9/22	9/23	9/24	9/25																		
平均 気温	本年値	27.5	27.6	28.0	28.1	28.5	27.9	25.9	26.5	27.6	29.0	29.9	28.1	22.9	23.2	20.5	25.6	27.1	28.3	27.9	23.7	22.3	22.9	25.4	24.5	24.0	21.1	21.2	21.7	23.7	25.6	24.1	21.5																												
	平年値	26.6	26.5	26.5	26.4	26.3	26.2	26.1	26.0	25.9	25.8	25.7	25.6	25.5	25.4	25.3	25.1	25.0	24.9	24.8	24.7	24.5	24.4	24.2	24.0	23.8	23.6	23.5	23.3	23.1	22.9	22.7	22.5	22.3	22.0	21.8	21.6	21.4	21.2	21.0	20.8	20.6	20.4																		
出 穂 期	7/10	992	1020	1048	1076	1104	1132	1158	1185	1,212	1,241																																																		
	7/11	965	993	1021	1049	1077	1105	1131	1158	1185	1,214	1,244																																																	
	7/12	938	965	993	1022	1050	1078	1104	1130	1158	1187	1,217	1,245																																																
	7/13	909	936	964	992	1021	1049	1075	1101	1129	1158	1188	1,216	1,239																																															
	7/14	881	908	936	964	993	1021	1047	1073	1101	1130	1160	1188	1,211	1,234																																														
	7/15	852	879	907	935	964	992	1018	1044	1072	1101	1131	1159	1182	1,205	1,225																																													
	7/16	824	852	880	908	936	964	990	1017	1044	1073	1103	1131	1154	1,177	1,198	1,223																																												
	7/17	797	825	853	881	909	937	963	990	1017	1046	1076	1104	1127	1,150	1,171	1,196	1,223																																											
	7/18		797	825	853	882	909	935	962	989	1018	1048	1076	1099	1,123	1,143	1,169	1,196	1,224																																										
	7/19			796	824	852	880	906	933	960	989	1019	1047	1070	1,093	1,114	1,139	1,167	1,195	1,223	1,246																																								
	7/20				794	822	850	876	903	930	959	989	1017	1040	1,063	1,084	1,109	1,137	1,165	1,193	1,216	1,239																																							
	7/21				765	793	821	847	874	901	930	960	988	1,011	1,034	1,055	1,080	1,108	1,136	1,164	1,187	1,210	1,233																																						
	7/22					792	818	844	872	901	931	959	982	1,005	1,025	1,051	1,078	1,106	1,134	1,158	1,180	1,203	1,228																																						
	7/23						789	815	843	872	902	930	953	976	996	1,022	1,049	1,077	1,105	1,129	1,151	1,174	1,200	1,224	1,248																																				
	7/24								814	843	873	901	924	947	968	993	1,020	1,049	1,076	1,100	1,122	1,145	1,171	1,195	1,219	1,240																																			
	7/25									816	846	874	897	920	940	966	993	1,021	1,049	1,073	1,095	1,118	1,143	1,168	1,192	1,213	1,234																																		
	7/26										821	849	872	895	916	942	969	997	1,025	1,049	1,071	1,094	1,119	1,144	1,168	1,189	1,210	1,232																																	
	7/27											824	847	870	891	916	943	972	1,000	1,023	1,046	1,068	1,094	1,118	1,142	1,163	1,185	1,206	1,230																																
	7/28												822	845	866	891	918	947	975	998	1,021	1,043	1,069	1,093	1,117	1,138	1,160	1,181	1,205	1,231																															
	7/29													820	840	866	893	921	949	973	995	1,018	1,043	1,068	1,092	1,113	1,134	1,156	1,180	1,205	1,229																														
	7/30														813	838	865	894	922	945	968	991	1,016	1,040	1,064	1,086	1,107	1,128	1,152	1,178	1,202	1,223	1,246																												
	7/31															810	837	866	894	917	940	963	988	1,012	1,036	1,058	1,079	1,100	1,124	1,150	1,174	1,195	1,218	1,240																											
	8/1																810	838	866	889	912	935	960	985	1,009	1,030	1,051	1,073	1,096	1,122	1,146	1,167	1,190	1,212	1,234																										
	8/2																	811	839	862	885	907	933	957	981	1,002	1,024	1,045	1,069	1,095	1,119	1,140	1,163	1,185	1,206	1,228	1,249																								
	8/3	収穫適期のめやす（出穂後の積算温度・通常年の場合） 【早生品種】 なつひめ　　・ ・ ・ ・ ・ 1, 000℃前後 五百万石、こしいぶき　　・ ・ ・ ・ 975℃ つきあかり　　・ ・ ・ ・ ・ 1, 000～1, 100℃ 【中生品種】 こがねもち、コシヒカリ　　・ ・ ・ 1, 000℃ 【晩生品種】 新之助　　・ ・ ・ ・ ・ 1, 050～1, 100℃ ○高温登熟条件となる場合には、積算温度を上記のめやすより50℃（2日程度） 早めましょう。 ○実際の収穫は粳の黄化割合で判断しましょう。																			811	835	857	880	905	930	954	975	996	1,018	1,041	1,067	1,091	1,113	1,135	1,157	1,179	1,200	1,222	1,243																					
8/4																																807	829	852	877	902	926	947	968	990	1,013	1,039	1,063	1,085	1,107	1,129	1,151	1,172	1,194	1,215	1,236										
8/5																																		801	824	849	874	898	919	940	962	985	1,011	1,035	1,056	1,079	1,101	1,123	1,144	1,166	1,187	1,208	1,229	1,249							
8/6																																				820	845	869	890	911	933	956	982	1,006	1,028	1,050	1,072	1,094	1,115	1,137	1,158	1,179	1,200	1,220	1,241						
8/7																																					815	839	860	881	903	927	952	976	998	1,020	1,042	1,064	1,086	1,107	1,128	1,149	1,170	1,191	1,211						
8/8																																												811	832	853	875	898	924	948	970	992	1,014	1,036	1,057	1,079	1,100	1,121	1,142	1,162	1,183
8/9																																													805	826	848	872	897	921	943	965	987	1,009	1,030	1,052	1,073	1,094	1,115	1,135	1,156
8/10																																														800	822	846	871	896	917	939	961	983	1,005	1,026	1,047	1,068	1,089	1,110	1,130
8/11																																																822	848	872	894	916	938	960	981	1,003	1,024	1,045	1,066	1,086	1,107
8/12																																																		822	846	867	889	911	933	955	976	997	1,018	1,039	1,060
8/13																																		818	840	862	884	906	928	949	970	991	1,012	1,033	1,053																
8/14																																			812	834	856	878	900	921	942	963	984	1,005	1,025																
8/15																																				807	829	850	872	893	915	936	956	977	997																